

2-2008

Graph-Based Upper Bounds for the Probability of the Union of Events

Pierangela Veneziani

The College at Brockport, pvenezia@brockport.edu

Follow this and additional works at: https://digitalcommons.brockport.edu/mth_facpub

 Part of the [Mathematics Commons](#)

Repository Citation

Veneziani, Pierangela, "Graph-Based Upper Bounds for the Probability of the Union of Events" (2008). *Mathematics Faculty Publications*. 8.

https://digitalcommons.brockport.edu/mth_facpub/8

Citation/Publisher Attribution:

Veneziani, P. (2008). Graph-Based Upper Bounds for the Probability of the Union of Events. *Electronic Journal of Combinatorics*, 15(1R), 1-11.

This Article is brought to you for free and open access by the Department of Mathematics at Digital Commons @Brockport. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of Digital Commons @Brockport. For more information, please contact kmyers@brockport.edu.

Graph-based upper bounds for the probability of the union of events

Pierangela Veneziani*

Mathematics Department
SUNY College at Brockport
350 New Campus Drive
Brockport, NY, US
pvenezia@brockport.edu

Submitted: Nov 27, 2007; Accepted: Jan 28, 2008; Published: Feb 11, 2008
Mathematics Subject Classifications: 60E15, 90C05

Abstract

We consider the problem of generating upper bounds for the probability of the union of events when the individual probabilities of the events as well as the probabilities of pairs of these events are known. By formulating the problem as a Linear Program, we can obtain bounds as objective function values corresponding to dual basic feasible solutions. The new upper bounds are based on underlying bipartite and threshold type graph structures.

1 Introduction

The *Boolean probability bounding problem* can be formulated as follows: let A_1, \dots, A_n be a finite set of arbitrary events in a probability space Ω , and let us assume that the individual probabilities $P(A_i)$, $i = 1, \dots, n$, as well as the probabilities $P\left(\bigcap_{1 \leq i_1 < \dots < i_l \leq n} A_{i_l}\right)$, $l = 2, \dots, m$, of up to m -tuples of these events are known, where $m < n$. Using this information we want to generate upper and lower bounds for the probability of a Boolean function of these events. The integer m is usually referred to as the *degree* of these bounds.

The Boolean probability bounding scheme is a particular instance of the optimization version of the *probabilistic satisfiability* (PSAT) *problem*. The decision version of PSAT consists in determining whether, given the probabilities that m logical sentences defined

*The author wishes to thank Dr. E. Boros for his continued support and her daughter Sofia for inspiration. The author would also like to acknowledge the work of the reviewer of the manuscript.

on n logical variables are true, such probability assignment is consistent. The optimization version of PSAT is concerned with determining bounds on the probability that an additional sentence is true. PSAT is known to be NP-hard (see e.g. [7]).

Both versions of PSAT were first proposed as general problem in the theory of probability by George Boole, who suggested algebraic methods to solve it.

Hailperin [5] formulated Boole's general problem as a Linear Program and showed that Boole's method is equivalent to Fourier's elimination. Kounias and Marin [8] utilized Hailperin's linear model in their work on the Boolean probability bounding scheme and generated bounds of degree two.

Dawson and Sankoff [4] proposed a sharp lower bound for the probability that at least one out of n events occurs, using the first two binomial moments of the occurrences and a linear programming formulation.

Hunter [6] found an upper bound of degree two expressed in terms of underlying spanning trees, Bukszár and Prékopa [2] by means of cherry-trees, and Bukszár and Szantai [3] by means of hypercherry trees.

Let us introduce the following notations: let $\mathcal{G}_m = (\mathcal{V}, \mathcal{E})$ denote the hypergraph where $\mathcal{V} = \{1, \dots, n\}$ and $\mathcal{E} = \bigcup_{k=2}^m \mathcal{E}_k$ where $\mathcal{E}_k = \{I \subseteq \mathcal{V} \mid |I| = k\}$, $k = 2, \dots, m$. Further let $\Gamma = \mathcal{V} \cup \mathcal{E}$.

For each subset $J \subseteq \mathcal{V}$ let us define the event $C_J = (\bigcap_{i \in J} A_i) (\bigcap_{i \in J^c} A_i^c)$ where $J^c = \mathcal{V} \setminus J$, and $A_i^c = \Omega \setminus A_i$, $i = 1, \dots, n$, and to each subset $J \subseteq \mathcal{V}$ let us associate a decision variable $x_J = \Pr(C_J)$ and a scalar c_J .

Let us further introduce the notation $p_I = P(\bigcap_{i \in I} A_i)$ where $I \in \Gamma$, and let us set $p_\emptyset = 1$ by definition.

Let us note that the equality $\sum_{I \subseteq J \subseteq \mathcal{V}} P(C_J) = P(\bigcap_{i \in I} A_i)$ holds for all subsets $I \in \Gamma \cup \{\emptyset\}$, because the 2^n (disjoint) events C_J 's form a partition of the probability space Ω . We can write this equivalently as $\sum_{I \subseteq J \subseteq \mathcal{V}} x_J = p_I$.

Finally, let p denote the vector with components $p_I \in [0, 1]$, $I \in \Gamma \cup \{\emptyset\}$, let x be the vector with components $x_J \in [0, 1]$, $J \subseteq \mathcal{V}$, and let $H = (h_{IJ})$ denote the incidence matrix whose entries are defined by $h_{IJ} = 1$ if $I \subseteq J$, $h_{IJ} = 0$ otherwise.

The matrix H has $\sum_{i=0}^m \binom{n}{i}$ rows and 2^n columns.

In the vectors p and x , and in the row and column indices of the matrix H the order of the elements will follow the lexicographic order of the subscript sets.

The Boolean probability bounding problem can thus be restated as a linear program of the form

$$\begin{aligned} & \text{Max or Min} \quad \sum_{J \subseteq \mathcal{V}} c_J x_J \\ & \text{st} \\ & \quad \sum_{I \subseteq J \subseteq \mathcal{V}} h_{IJ} x_J = p_I \quad \forall I \in \Gamma \cup \{\emptyset\} \\ & \quad x_J \geq 0 \quad \forall J \subseteq \mathcal{V} \end{aligned}$$

larger than 1 then, by taking into account the constraint $\sum_{J \subseteq \mathcal{V}} x_J = 1$, we can trivially set the upper bound to 1. Therefore the first row of the matrix H as well as the first column corresponding to the variable x_\emptyset can be disregarded from our formulation. In (1) we now have $n + \binom{n}{2}$ constraints and $2^n - 1$ variables.

As Prékopa et al. suggested in [11], it is then possible to interpret the $n + \binom{n}{2}$ components of any dual feasible solution $w = (w_\gamma)_{\gamma \in \Gamma}$ of problem (2) as node and edge weights in \mathcal{G}_2 , that is a weight w_i is assigned to node $i \in \mathcal{V}$ and a weight $w_{i,j}$ is assigned to edge $\{i, j\} \in \mathcal{E}_2$.

In what follows we will let $\mathcal{E}(S)$ denote the edge set of a subset $S \subseteq \mathcal{V}$ and $w(S) = \sum_{\gamma \in S} w_\gamma + \sum_{\gamma \in \mathcal{E}(S)} w_\gamma$ represent the weight of subset S for a given dual feasible solution $w = (w_\gamma)_{\gamma \in \Gamma}$.

For the instance under study ($c^T = [0, 1, \dots, 1]$ and $m = 2$) problem (2) can then be written as

$$\begin{aligned} \text{Min} \quad & \sum_{\gamma \in \Gamma} p_\gamma w_\gamma \\ \text{st} \quad & w(S) \geq 1 \quad \forall S \subseteq \mathcal{V}. \end{aligned} \tag{3}$$

The lemma that follows provides a sufficient and necessary condition for a given vector to be a basic feasible solution of problem (3) by use of the graph structure introduced at the beginning of section 2.

Lemma 1 *Given a collection $\mathfrak{S} = \{I_\gamma\}_{\gamma \in \Gamma}$ of column subscripts of the matrix H , a vector $w = (w_\gamma)_{\gamma \in \Gamma}$ is a dual basic feasible solution of problem (1) generated by the basis \mathfrak{S} if the following conditions are satisfied.*

(i) *The vector $w = (w_\gamma)_{\gamma \in \Gamma}$ is the unique solution of the system of equations $w(I_\gamma) = 1$ for all subsets $I_\gamma \in \mathfrak{S}$, $\gamma \in \Gamma$.*

(ii) *For all subsets $S \subseteq \mathcal{V}$ such that $S \notin \mathfrak{S}$ the inequality $w(S) \geq 1$ holds.*

Proof. Let h_J , $J \subseteq \mathcal{V}$, designate a column vector of the matrix H . Let B denote a nonsingular square submatrix of H of order m and let $\mathfrak{S} = \{I_\gamma\}_{\gamma \in \Gamma}$ denote the collection of subscripts whose columns form B . Recall that a matrix B is said to be a dual feasible basis of problem (1) if $c_B^T B^{-1} h_{I_\gamma} = c_{I_\gamma}$ for all subsets $I_\gamma \in \mathfrak{S}$, $\gamma \in \Gamma$, and $c_B^T B^{-1} h_J \geq c_J$ for all subsets $J \notin \mathfrak{S}$. The corresponding dual basic feasible solution is the vector $w^T = c_B^T B^{-1}$.

In our case, condition (i) guarantees that the matrix B is nonsingular and that the equalities $c_B^T B^{-1} h_{I_\gamma} = c_{I_\gamma}$ hold for all basic sets $I_\gamma \in \mathfrak{S}$, $\gamma \in \Gamma$, and condition (ii) ensures that the inequalities $c_B^T B^{-1} h_J \geq c_J$ are satisfied for all nonbasic sets $J \notin \mathfrak{S}$. ■

Remark 2 *Let $\mathcal{G}^* = (\Gamma \cup \mathfrak{S}, \mathcal{E}^*)$ denote the bipartite graph where $\mathcal{E}^* = \{I \in \Gamma, J \in \mathfrak{S} \mid I \subseteq J\}$. A necessary condition for a collection $\mathfrak{S} = \{I_\gamma\}_{\gamma \in \Gamma}$ of column subscripts of H to form a basis is that there exists a perfect matching in the bipartite graph \mathcal{G}^* , otherwise if no perfect matching exists the matrix B would be singular (see e.g. [10]). Therefore in constructing a basis $\mathfrak{S} = \{I_\gamma\}_{\gamma \in \Gamma}$ we want to make sure to cover all the nodes and edges of \mathcal{G}_2 .*

The new families of dual feasible bases presented in the propositions that follow are obtained by partitioning the graph \mathcal{G}_2 in two components and then assigning nonnegative weights to each component and nonpositive weights to the edges connecting them.

Proposition 3 Assume $n \geq 5$. Let $\mathcal{V} = \mathcal{A} \cup \mathcal{B}$ be a partition of the vertex set \mathcal{V} and $k = |\mathcal{A}|$ where $2 \leq k \leq \lfloor \frac{n}{2} \rfloor$. Let l denote a positive integer such that $k \leq l \leq n - k - 1$, $\binom{n-k}{2} \leq \binom{n-k}{l}$, and $\binom{k}{2} \leq \binom{n-k}{l+1}$. Then the vector $w = (w_\gamma)_{\gamma \in \Gamma}$ with components

$$w_\gamma = \begin{cases} 1 & \text{if } \gamma \in \mathcal{V} \\ -1 & \text{if } \gamma = \{i, j\}, i \in \mathcal{A}, j \in \mathcal{B} \\ \frac{l-1}{k} & \text{if } \gamma = \{i, j\}, i \in \mathcal{A}, j \in \mathcal{A} \\ \frac{k-1}{l} & \text{if } \gamma = \{i, j\}, i \in \mathcal{B}, j \in \mathcal{B} \end{cases}$$

is a dual basic feasible solution of problem (1).

Proof. Let $\mathcal{A}_s \subseteq \mathcal{A}$ ($\mathcal{B}_r \subseteq \mathcal{B}$) denote a subset of cardinality s , $1 \leq s \leq k$ (r , $1 \leq r \leq n - k$).

Define $\mathfrak{S} = \{I_\gamma\}_{\gamma \in \Gamma}$ to be the collection of column labels of the matrix H where

$$I_i = \{i\} \text{ for } i \in \mathcal{V}$$

$$I_{i,j} = \begin{cases} \{i, j\} & \text{if } i \in \mathcal{A}, j \in \mathcal{B} \\ \mathcal{A} \cup \mathcal{B}_{l+1} & \text{if } i, j \in \mathcal{A} \\ \mathcal{A} \cup \mathcal{B}_l & \text{if } i, j \in \mathcal{B}, \text{ where } i, j \in \mathcal{B}_l. \end{cases}$$

The vector w is a dual basic feasible solution of problem (1) generated by the basis \mathfrak{S} because conditions (i) and (ii) of lemma 1 are met, as shown below.

(i) For all $i \in \mathcal{V}$ $w(I_i) = 1$ if and only if $w_i = 1$.

For all $\{i, j\} \in \mathcal{E}_2$ such that $i \in \mathcal{A}$, $j \in \mathcal{B}$ $w(I_{i,j}) = w(\{i, j\}) = w_i + w_j + w_{i,j} = 2 + w_{i,j} = 1$ if and only if $w_{i,j} = -1$.

The symmetry of the constraints ensures that $w_{h,k} = x$ for all $h, k \in \mathcal{A}$ and $w_{h,k} = y$ for all $h, k \in \mathcal{B}$.

Then for all $\{i, j\} \in \mathcal{E}(\mathcal{A})$

$$\begin{aligned} w(I_{i,j}) = w(\mathcal{A} \cup \mathcal{B}_{l+1}) &= \sum_{f \in \mathcal{A} \cup \mathcal{B}_{l+1}} w_f + \sum_{f,g \in \mathcal{A}} w_{f,g} + \sum_{f,g \in \mathcal{B}_{l+1}} w_{f,g} + \sum_{f \in \mathcal{A}, g \in \mathcal{B}_{l+1}} w_{f,g} \\ &= (k + l + 1)(1) + \binom{k}{2}x + \binom{l+1}{2}y + k(l+1)(-1) = 1 \end{aligned}$$

if and only if

$$\frac{k(k-1)}{2}x + \frac{(l+1)l}{2}y = kl - l. \tag{I}$$

For all $\{i, j\} \in \mathcal{E}(\mathcal{B})$

$$\begin{aligned} w(I_{i,j}) = w(\mathcal{A} \cup \mathcal{B}_l) &= \sum_{f \in \mathcal{A} \cup \mathcal{B}_l} w_f + \sum_{f,g \in \mathcal{A}} w_{f,g} + \sum_{f,g \in \mathcal{B}_l} w_{f,g} + \sum_{f \in \mathcal{A}, g \in \mathcal{B}_l} w_{f,g} \\ &= (k + l)(1) + \binom{k}{2}x + \binom{l}{2}y + kl(-1) = 1 \end{aligned}$$

if and only if

$$\frac{k(k-1)}{2}x + \frac{l(l-1)}{2}y = kl - k - l + 1. \quad (\text{II})$$

The unique solution of the system of equations $I - II$ is given by $x = \frac{l-1}{k}$ and $y = \frac{k-1}{l}$.

(ii) The only nontrivial case that needs to be considered to prove that the vector w is feasible for problem (3) is $S = \mathcal{A}_s \cup \mathcal{B}_r$, $1 \leq s \leq k$, $1 \leq r \leq n - k$.

Then

$$\begin{aligned} w(S) = w(\mathcal{A}_s \cup \mathcal{B}_r) &= \sum_{f \in \mathcal{A}_s \cup \mathcal{B}_r} w_f + \sum_{f, g \in \mathcal{A}_s} w_{f, g} + \sum_{f, g \in \mathcal{B}_r} w_{f, g} + \sum_{f \in \mathcal{A}_s, g \in \mathcal{B}_r} w_{f, g} \\ &= s + r + \binom{s}{2}x + \binom{r}{2}y + rs(-1) \\ &= \frac{yr^2 + r(2 - 2s - y) + xs^2 - xs + 2s}{2}. \end{aligned}$$

Consider the following cases.

Case 1: $s = 1$. Then $w(\mathcal{A}_1 \cup \mathcal{B}_r) = 1 + \frac{yr(r-1)}{2} \geq 1$ because $r \geq 1$ and $y \geq 0$.

Case 2: $s = k$. Then $w(\mathcal{A} \cup \mathcal{B}_r) = \frac{r^2 - r(1+2l) + l(l+1) + 2}{2}$. Therefore the inequality $w(\mathcal{A} \cup \mathcal{B}_r) \geq 1$ holds if and only if $r^2 - r(1 + 2l) + l(l + 1) \geq 0$ equivalently $r \leq l$ or $r \geq l + 1$.

Case 3: $2 \leq s \leq k - 1$. Consider the polynomial

$$\mathcal{P}(r|s, l) = w(S) - 1 = \frac{yr^2 + r(2 - 2s - y) + xs^2 - xs + 2s}{2} - 1$$

where $r \in [1, n - k]$, and let $\Delta(s, l)$ denote the discriminant of the equation $\mathcal{P}(r|s, l) = 0$.

We will show that $\Delta(s, l) < 0$ for every $s \in [2, k - 1]$, $l \in [k, n - k - 1]$, thus $\mathcal{P}(r|s, l) > 0$ for any r , equivalently $w(\mathcal{A}_s \cup \mathcal{B}_r) \geq 1$ for any subset $\mathcal{A}_s \cup \mathcal{B}_r \subseteq \mathcal{V}$.

The expression of the discriminant is given by

$$\begin{aligned} \Delta(s, l) &= (2 - 2s - y)^2 - 4y(xs^2 - xs + 2s - 2) \\ &= \frac{4(1 - s)(k - s)l^2 + 4(k - 1)(1 - s)(k - s)l + k(k - 1)^2}{kl^2}. \end{aligned}$$

The inequality $\Delta(s, l) < 0$ can be written as $\Delta^*(s, l) = -kl^2\Delta(s, l) > 0$.

Because the partial derivative of the function $\Delta^*(s, l)$ with respect to the variable l

$$\frac{\partial \Delta^*(s, l)}{\partial l} = 8(s - 1)(k - s)l + 4(k - 1)(s - 1)(k - s) = 4(s - 1)(k - s)(2l + k - 1)$$

is positive when $l \geq -(\frac{k-1}{2})$, the function $\Delta^*(s, l)$ is increasing in l on the interval $[k, n - k - 1]$. If we show that the inequality $\Delta^*(s, l = k) \geq 0$ is satisfied for any $s \in [2, k - 1]$, the argument will be complete.

Because $\Delta^*(s, l = k) = k [4(1 - 2k)s^2 + 4(2k^2 + k - 1)s - (3k - 1)^2]$, the sign of the first derivative $\frac{d(\Delta^*(s, l=k))}{ds} = k [8(1 - 2k)s + 4(2k^2 + k - 1)]$ is nonnegative for $s \leq 4(k+1)$, that is the function $\Delta^*(s, l = k)$ is increasing in s on the interval $[2, k - 1]$.

Therefore to conclude that the function $\Delta^*(s, l = k)$ is nonnegative on the interval $s \in [2, k - 1]$ it then suffices to notice that $\Delta^*(s = 2, l = k) = 7k^2 - 18k + 7 \geq 13 > 0$, because in the case under study $k \geq 3$. ■

The bound generated by evaluating the objective function of problem (3) at the dual basic feasible solution described by the above proposition is given by

$$P\left(\bigcup_{i=1}^n A_i\right) \leq \sum_{j \in \mathcal{V}} p_j - \sum_{i \in \mathcal{A}, j \in \mathcal{B}} p_{i,j} + \frac{l-1}{k} \sum_{i,j \in \mathcal{A}} p_{i,j} + \frac{k-1}{l} \sum_{i,j \in \mathcal{B}} p_{i,j}.$$

Remark 4 *The assumptions $k \leq l \leq n - k - 1$, $\binom{n-k}{2} \leq \binom{n-k}{l}$, and $\binom{k}{2} \leq \binom{n-k}{l+1}$ guarantee that there is a sufficient number of sets needed to form a basis as described in the above proof. Moreover $l = k$ is the smallest value for which the proposition holds true.*

Proposition 5 *Assume $n \geq 5$. Let $\mathcal{V} = \mathcal{A} \cup \mathcal{B}$ be a partition of the vertex set \mathcal{V} and $l = |\mathcal{A}|$ where $1 \leq l \leq n - 1$. Then the vector $w = (w_\gamma)_{\gamma \in \Gamma}$ with components*

$$w_\gamma = \begin{cases} 1 & \text{if } \gamma \in \mathcal{V} \\ -1 & \text{if } \gamma = \{i, j\}, i \in \mathcal{A}, j \in \mathcal{B} \\ 0 & \text{if } \gamma = \{i, j\}, i, j \in \mathcal{A} \\ l-1 & \text{if } \gamma = \{i, j\}, i, j \in \mathcal{B} \end{cases}$$

is a dual basic feasible solution of problem (1).

Proof. Define $\mathfrak{S} = \{I_\gamma\}_{\gamma \in \Gamma}$ to be the collection of column labels of the matrix H where

$$I_i = \{i\} \text{ for } i \in \mathcal{V}$$

$$I_{i,j} = \begin{cases} \{i\} \cup \{j\} & \text{if } i \in \mathcal{A}, j \in \mathcal{B} \\ \{i\} \cup \{j\} \cup \{k\} & \text{if } i, j \in \mathcal{A}, \text{ where } k \in \mathcal{B} \\ \{i\} \cup \{j\} \cup \mathcal{A} & \text{if } i, j \in \mathcal{B} \end{cases}$$

The vector w is a dual basic feasible solution of problem (1) generated by the basis \mathfrak{S} because conditions (i) and (ii) of lemma 1 are met, as shown below.

(i) For all $i \in \mathcal{V}$ $w(I_i) = 1$ if and only if $w_i = 1$.

For all $\{i, j\} \in \mathcal{E}_2$ with $i \in \mathcal{A}, j \in \mathcal{B}$ $w(I_{i,j}) = w(\{i\} \cup \{j\}) = w_i + w_j + w_{i,j} = 2 + w_{i,j} = 1$ if and only if $w_{i,j} = -1$.

For all $\{i, j\} \in \mathcal{E}(\mathcal{A})$

$$w(I_{i,j}) = w(\{i\} \cup \{j\} \cup \mathcal{A}) = w_i + w_j + w_k + w_{i,j} + w_{i,k} + w_{j,k} = 3 - 2 + w_{i,j} = 1$$

if and only if $w_{i,j} = 0$.

Finally for all $i, j \in \mathcal{E}(\mathcal{B})$

$$\begin{aligned} w(I_{i,j}) = w(\{i\} \cup \{j\} \cup \mathcal{A}) &= \sum_{h \in \{i\} \cup \{j\} \cup \mathcal{A}} w_h + \sum_{h \in \mathcal{A}} w_{h,i} + \sum_{h \in \mathcal{A}} w_{h,j} + \sum_{h,k \in \mathcal{A}} w_{h,k} + w_{i,j} \\ &= l + 2 + 2l(-1) + \binom{l}{2}(0) + w_{i,j} \\ &= 2 - l + w_{i,j} = 1 \end{aligned}$$

if and only if $w_{i,j} = l - 1$.

(ii) The only nontrivial case that need to be considered to prove that the vector w is feasible for problem (3) is $S = \mathcal{A}_s \cup \mathcal{B}_r$, where $\mathcal{A}_s \subseteq \mathcal{A}$, $|\mathcal{A}_s| = s$, $1 \leq s \leq l$, and $\mathcal{B}_r \subseteq \mathcal{B}$, $|\mathcal{B}_r| = r$, $1 \leq r \leq n - l$. Then

$$\begin{aligned} w(\mathcal{A}_s \cup \mathcal{B}_r) &= \sum_{f \in \mathcal{A}_s \cup \mathcal{B}_r} w_f + \sum_{f,g \in \mathcal{A}_s} w_{f,g} + \sum_{f,g \in \mathcal{B}_r} w_{f,g} + \sum_{f \in \mathcal{A}_s, g \in \mathcal{B}_r} w_{f,g} \\ &= s + r + (l - 1) \binom{r}{2} - sr \\ &= \frac{(l - 1)r^2 - r(2s - 2 + l - 1) + 2s}{2}. \end{aligned}$$

Therefore the inequality $w(\mathcal{A}_s \cup \mathcal{B}_r) \geq 1$ holds if and only if $(l - 1)r^2 - r(2s - 2 + l - 1) + 2s - 2 \geq 0$ equivalently $r \leq \frac{s-2}{l-1}$ or $r \geq 1$, since $\frac{s-2}{l-1} < 1$ because $s < l + 1$. ■

The bound generated by evaluating the objective function of problem (3) at the basic feasible solution described by the above proposition is given by

$$P\left(\bigcup_{i=1}^n A_i\right) \leq \sum_{i \in \mathcal{V}} p_i + (l - 1) \sum_{i,j \in \mathcal{B}} p_{i,j} - \sum_{i \in \mathcal{A}, j \in \mathcal{B}} p_{i,j}.$$

The following corollary shows that a known bound [11] can be obtained as a special case of the bound presented in proposition 7.

Corollary 6 Assume $n \geq 5$. For fixed $i_1, i_2 \in \mathcal{V}$, let $\mathcal{A} = \mathcal{V} \setminus \{\{i_1\} \cup \{i_2\}\}$. Then the vector $w = (w_\gamma)_{\gamma \in \Gamma}$ with components

$$w_\gamma = \begin{cases} 1 & \text{if } \gamma \in \mathcal{V} \\ -1 & \text{if } \gamma = \{i_k, j\}, k = 1, 2, j \in \mathcal{A} \\ 0 & \text{if } \gamma = \{i, j\}, i, j \in \mathcal{A} \\ n - 3 & \text{if } \gamma = \{i_1, i_2\} \end{cases}$$

is a dual basic feasible solution of problem (1).

Proof. Set $\mathcal{B} = \{i_1, i_2\}$ in proposition 6. ■

We conclude the section presenting a new bound that is obtained by means of an underlying threshold-type graph structure.

Proposition 7 Assume $n \geq 5$. Let $\mathcal{V} = \mathcal{A} \cup \mathcal{B}$ be a partition of the vertex set \mathcal{V} . Let $s = |\mathcal{A}|$, $1 \leq s \leq n - 1$, and $\mathcal{A} = \{a_1, a_2, \dots, a_s\}$.

Let $\mathcal{N}(a_k)$, $k = 1, \dots, s$, denote the set of vertices $i \in \mathcal{B}$ that are connected to vertex a_k . Assume that $\mathcal{N}(a_1) = \mathcal{B}$, $\mathcal{N}(a_h) \subseteq \mathcal{N}(a_k)$ if $h > k$, and $\mathcal{N}(a_h) \cap \mathcal{N}(a_k) \neq \emptyset$ for all $h, k = 1, \dots, s$. Then the vector $w = (w_\gamma)_{\gamma \in \Gamma}$ with component

$$w_\gamma = \begin{cases} 1 & \text{if } \gamma \in \mathcal{V} \\ -1 & \text{if } \gamma = \{a_k, j\}, j \in \mathcal{N}(a_k), 1 \leq k \leq s \\ |\mathcal{N}(a_h) \cap \mathcal{N}(a_k)| - 1 & \text{if } \gamma = \{a_h, a_k\}, 1 \leq h, k \leq s \\ 0 & \text{otherwise} \end{cases}$$

is a dual basic feasible solution of problem (1).

Proof. Define $\mathfrak{S} = \{I_\gamma\}_{\gamma \in \Gamma}$ to be the collection of column labels of the matrix H where

$$I_i = \{i\} \quad \text{for } i \in \mathcal{V}$$

$$I_{i,j} = \begin{cases} \{a_k, j\} & \text{if } i = a_k, j \in \mathcal{N}(a_k), 1 \leq k \leq s \\ \{\mathcal{N}(a_h) \cap \mathcal{N}(a_k)\} \cup \{a_h\} \cup \{a_k\} & \text{if } i = a_h, j = a_k, 1 \leq h, k \leq s \\ \{a_1\} \cup \{i\} \cup \{j\} & \text{if } i, j \in \mathcal{B} \\ \{a_h\} \cup \{a_k\} \cup \{j\} \cup \mathcal{N}(a_k) & \text{if } i = a_k, j \in \mathcal{B} \setminus \mathcal{N}(a_k), 1 \leq k \leq s \\ & \text{where } h = \max\{t \mid j \in \mathcal{N}(a_t)\}. \end{cases}$$

The vector w is a dual basic feasible solution of problem (1) generated by the basis \mathfrak{S} because conditions (i) and (ii) of lemma 1 are met, as shown below.

(i) For all $i \in \mathcal{V}$ $w(I_i) = 1$ if and only if $w_i = 1$.

For all $\{a_k, j\} \in \mathcal{E}_2$ with $j \in \mathcal{N}(a_k)$, $1 \leq k \leq s$,

$$w(I_{a_k,j}) = w(\{a_k, j\}) = w_{a_k} + w_j + w_{a_k,j} = 2 + w_{a_k,j} = 1$$

if and only if $w_{a_k,j} = -1$.

For all $\{i, j\} \in \mathcal{E}(\mathcal{B})$

$$w(I_{i,j}) = w(\{a_1\} \cup \{i\} \cup \{j\}) = w_{a_1} + w_i + w_j + w_{a_1,i} + w_{a_1,j} + w_{i,j} = 3 - 2 + w_{i,j} = 1$$

if and only if $w_{i,j} = 0$.

For $\{i, j\} = \{a_h, a_k\}$, $1 \leq h, k \leq s$,

$$\begin{aligned} w(I_{a_h,a_k}) &= w(\{\mathcal{N}(a_h) \cap \mathcal{N}(a_k)\} \cup \{a_h\} \cup \{a_k\}) \\ &= \sum_{f \in \{\mathcal{N}(a_h) \cap \mathcal{N}(a_k)\} \cup \{a_h\} \cup \{a_k\}} w_f + \sum_{f \in \mathcal{N}(a_h) \cap \mathcal{N}(a_k)} w_{a_h,f} + \sum_{f \in \mathcal{N}(a_h) \cap \mathcal{N}(a_k)} w_{a_k,f} \\ &\quad + \sum_{f,g \in \mathcal{N}(a_h) \cap \mathcal{N}(a_k)} w_{f,g} + w_{a_h,a_k} \\ &= |\mathcal{N}(a_h) \cap \mathcal{N}(a_k)| + 2 + |\mathcal{N}(a_h) \cap \mathcal{N}(a_k)|(-1) + w_{a_h,a_k} \\ &\quad + |\mathcal{N}(a_h) \cap \mathcal{N}(a_k)|(-1) + \binom{|\mathcal{N}(a_h) \cap \mathcal{N}(a_k)|}{2}(0) \\ &= 2 - |\mathcal{N}(a_h) \cap \mathcal{N}(a_k)| + w_{a_h,a_k} = 1 \end{aligned}$$

if and only if $w_{a_h,a_k} = |\mathcal{N}(a_h) \cap \mathcal{N}(a_k)| - 1$.

Finally for $\{a_k, j\} \in \mathcal{E}_2$ with $j \in \mathcal{B} \setminus \mathcal{N}(a_k)$, $1 \leq k \leq s$,

$$\begin{aligned} w(I_{a_k,j}) &= w(\{a_h\} \cup \{a_k\} \cup \{j\} \cup \mathcal{N}(a_k)) \\ &= w_{a_h} + w_{a_k} + w_j + \sum_{f \in \mathcal{N}(a_k)} w_f + w_{a_h,a_k} + w_{a_h,j} + w_{a_k,j} \\ &\quad + \sum_{f \in \mathcal{N}(a_k)} w_{a_h,f} + \sum_{f \in \mathcal{N}(a_k)} w_{a_k,f} + \sum_{f \in \mathcal{N}(a_k)} w_{j,f} \\ &= 3 + |\mathcal{N}(a_k)| + |\mathcal{N}(a_h) \cap \mathcal{N}(a_k)| - 1 - 1 + w_{a_k,j} \\ &\quad + |\mathcal{N}(a_k)|(-1) + |\mathcal{N}(a_k)|(-1) + |\mathcal{N}(a_k)|(0) \\ &= 1 + w_{a_k,j} = 1 \end{aligned}$$

if and only if $w_{a_k,j} = 0$, since $|\mathcal{N}(a_h) \cap \mathcal{N}(a_k)| = |\mathcal{N}(a_k)|$ because $k > h$.

(ii) The only nontrivial case that needs to be considered to prove that the vector w is feasible for problem (2) is $S = C \cup D$, where $C \subseteq \mathcal{A}$ and $D \subseteq \mathcal{B}$. We will show that the inequality $w(C \cup D) \geq 1$ holds by induction on $|C|$.

If $|C| = 1$ let $C = \{a_{i_1}\}$, where $1 \leq i_1 \leq s$. Then

$$\begin{aligned} w(S) &= w(\{a_{i_1}\} \cup D) = \sum_{j \in \{a_{i_1}\} \cup D} w_j + \sum_{j \in D} w_{a_{i_1},j} + \sum_{i,j \in D} w_{i,j} \\ &= 1 + |D| + |\mathcal{N}(a_{i_1}) \cap D|(-1) + 0 \geq 1 \end{aligned}$$

because $|\mathcal{N}(a_{i_1}) \cap D| \leq |D|$.

Let us assume that inequality $w(C \cup D) \geq 1$ holds for $|C| = k$.

If $|C| = k + 1$ let $C = \{a_{i_1}, a_{i_2}, \dots, a_{i_k}, a_{i_{k+1}}\}$, where $1 \leq i_1 < i_2 < \dots < i_k < i_{k+1} \leq s$, then

$$\begin{aligned} w(S) &= w(\{a_{i_1}, a_{i_2}, \dots, a_{i_{k+1}}\} \cup D) \\ &= w(\{a_{i_1}, a_{i_2}, \dots, a_{i_k}\} \cup D) + w_{a_{i_{k+1}}} + \sum_{j \in D} w_{a_{i_{k+1}},j} + \sum_{l=1}^k w_{a_{i_l}, a_{i_{k+1}}} \\ &= w(\{a_{i_1}, a_{i_2}, \dots, a_{i_k}\} \cup D) + 1 + |\mathcal{N}(a_{i_{k+1}}) \cap D|(-1) \\ &\quad + \sum_{l=1}^k [|\mathcal{N}(a_{i_{k+1}}) \cap \mathcal{N}(a_{i_l})| - 1] \\ &= w(\{a_{i_1}, a_{i_2}, \dots, a_{i_k}\} \cup D) + 1 - |\mathcal{N}(a_{i_{k+1}}) \cap D| + |\mathcal{N}(a_{i_{k+1}}) \cap \mathcal{N}(a_{i_1})| - 1 \\ &\quad + \sum_{l=2}^k [|\mathcal{N}(a_{i_{k+1}}) \cap \mathcal{N}(a_{i_l})| - 1] \\ &= w(\{a_{i_1}, a_{i_2}, \dots, a_{i_k}\} \cup D) + |\mathcal{N}(a_{i_{k+1}}) \cap \mathcal{N}(a_{i_1})| \\ &\quad - |\mathcal{N}(a_{i_{k+1}}) \cap D| + \sum_{l=2}^k [|\mathcal{N}(a_{i_{k+1}}) \cap \mathcal{N}(a_{i_l})| - 1] \geq 1 \end{aligned}$$

because $w(\{a_{i_1}, a_{i_2}, \dots, a_{i_k}\} \cup D) \geq 1$ by the induction hypothesis and $|\mathcal{N}(a_{i_{k+1}}) \cap \mathcal{N}(a_{i_1})| - |\mathcal{N}(a_{i_{k+1}}) \cap D| \geq 0$ since $|\mathcal{N}(a_{i_{k+1}}) \cap \mathcal{N}(a_{i_1})| = |\mathcal{N}(a_{i_{k+1}})| \geq |\mathcal{N}(a_{i_{k+1}}) \cap D|$. ■

The bound generated by evaluating the objective function of problem (3) at the dual basic feasible solution described by the above proposition is given by

$$P\left(\bigcup_{i=1}^n A_i\right) \leq \sum_{i \in \mathcal{V}} p_i + \sum_{\substack{h,k=1 \\ h \neq k}}^s [|\mathcal{N}(a_h) \cap \mathcal{N}(a_k)| - 1] p_{a_h, a_k} - \sum_{k=1}^s \sum_{j \in \mathcal{N}(a_k)} p_{a_k, j}.$$

Finally we will compare one of the new bounds with Kwerel's bound [9] and Hunter's bound [6] for the system II in [1], for which $n = 6$, $P\left(\bigcup_{i=1}^n A_i\right) = 0.6740$,

$p_1 = 0.268, p_2 = 0.312, p_3 = 0.302, p_4 = 0.172, p_5 = 0.384, p_6 = 0.278,$
 $p_{12} = 0, p_{13} = 0.168, p_{14} = 0.033, p_{15} = 0.19, p_{16} = 0.155,$
 $p_{23} = 0.078, p_{24} = 0.045, p_{25} = 0.156, p_{26} = 0.067,$
 $p_{34} = 0.056, p_{35} = 0.201, p_{36} = 0.111,$
 $p_{45} = 0.049, p_{46} = 0.089, p_{56} = 0.189.$

Numerical results for the given system are given in the table below.

Upper Bounds of Degree 2	
<i>Kwerel's bound</i>	1
<i>Hunter's bound</i>	0.891
<i>Our bound*</i>	0.813
<i>Exact value</i>	0.674

*Our bound was generated by setting $\mathcal{A} = \{1, 2, 4\}$, $\mathcal{B} = \{3, 5, 6\}$, and $l = 1$ in proposition 6. It is possible to show that our bound is the best possible bound that can be generated via the linear programming formulation of the Boolean Probability Bounding Problem for the numerical example under study.

References

- [1] F. Alajaji, H. Kuai and G. Takahara, A Lower Bound for the Probability of a Finite Union of Events *Discrete Applied Mathematics*, **215** (2000), 147 - 158.
- [2] J. Bukszár and A. Prékopa, Probability bounds with cherry trees, *Math. Oper. Res.*, **26** (2001), 174 - 192.
- [3] J. Bukszár and T. Szántai, Probability bounds given by hypercherry trees, *Alkalmaz. Mat. Lapok*, **19** (1999), 69 - 85.
- [4] D.A. Dawson and S. Sankoff, An Inequality for Probability, . *Proc. Am. Math. Soc.*, **18** (1967), 504-507.
- [5] T. Hailperin, Best possible inequalities for the probability of a logical function of events, *The American Mathematical Monthly*, **72** (1965), 343 - 359.
- [6] D. Hunter, An upper bound for the probability of the union, *J. Appl. Prob.*, **30** (1975), 597 - 603.
- [7] B. Jaumard, P. Hansen and M. Poggi de Aragão, Column generation methods for probabilistic logic, *ORSA J. Comput.*, **3** (1991), 135 - 148.
- [8] S. Kounias and J. Marin, Best linear bonferroni bounds, *SIAM J. on Applied Mathematics*, **30** (1976), 301 - 326.
- [9] S.M. Kwerel, Most stringent bounds on aggregated probabilities of partially specified dependent probability systems, *J. Am. Statist. Assoc.* **70** (1975) 472-479.
- [10] L. Lovász and M.D. Plummer, Matching theory, *Ann. Discrete Math.*, **29** (1986).
- [11] A. Prékopa, B. Vizvári and G. Regös, Lower and upper bounds on probabilities of Boolean functions of events, *Rutcor Research Report*, **36-95** (1995)